

Instructions for completing Medical Exemption Form:								
Section 1: Completed by parent/guardian or student (aged >18 years): Enter child care facility, school, or post-secondary school, and student information								
	Completed by licensed health care provider (MD, DO, ND, APRN-Rx, PA): Check exempted vaccine, contraindication or precaution, or both, and							
	mplete duration of exemption							
Section 1: Child Care Facility, School, Post-Secondary School, and Student Information								
Student's Name:		Student's Date of Birth:						
Student's Home Address		City	State	Zip				
Name of Child Care Facility, School, Post-Secondary School		Street Address	City	Zip				
			,	•				
I understand that if at any time there is, in the opinion of the Department of Health, danger of an outbreak or epidemic from any communicable disease for which								
	required, this exemption from immunization shall not be recogniz							
	facility, school, or post-secondary school until the Director of Health has determined that the presence of the outbreak no longer exists [HRS §302A-1157].							
Parent/Guardia	n Name [if student <18 years]. (Please print):							
Parent/Guardia	an <b>or</b> Student (if aged <u>&gt;</u> 18 years) Signature:	Date	2:					
Section 2: Fo	r Health Care Provider Use ONLY (MD, DO, ND, APRN-Rx	. PA):						
VACCINE	<b>CONTRAINDICATIONS*</b> (Check all that apply to this patient):	<b>PRECAUTIONS*</b> (Check all that apply to this patient)	FROM:	To:				
	Severe allergic reaction (e.g., anaphylaxis) after a	□ Guillain-Barre Syndrome <6 weeks after previous	/ /	/ /				
	previous dose or to a vaccine component			/ /				
		dose of tetanus-toxoid-containing vaccine	/ /					
🗆 Tdan		dose of tetanus-toxoid-containing vaccine History of Arthus-type hypersensitivity reactions						
🗆 Tdap	<b>DTaP/Tdap only</b> : Encephalopathy (e.g., coma,	History of Arthus-type hypersensitivity reactions						
-	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged	History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-						
<ul> <li>Tdap</li> <li>DT, Td</li> </ul>	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable	History of Arthus-type hypersensitivity reactions						
-	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid- containing or tetanus-toxoid-containing vaccine</li> </ul>						
-	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid- containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without</li> </ul>						
-	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid- containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> </ul>						
-	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable</li> </ul>						
-	DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms,</li> </ul>						
DT, Td	<ul> <li>DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, Tdap</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> </ul>	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy</li> </ul>		/ /				
DT, Td	<ul> <li>DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, Tdap</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Age &lt;6 weeks</li> </ul>	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy</li> <li>Moderate or severe acute illness with or without fever</li> </ul>		/ /				
DT, Td	<ul> <li>DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, Tdap</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> </ul>	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy</li> <li>Moderate or severe acute illness with or without</li> </ul>		/ /				
<ul> <li>DT, Td</li> <li>Hib</li> <li>Hep A</li> </ul>	<ul> <li>DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, Tdap</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Age &lt;6 weeks</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> </ul>	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy</li> <li>Moderate or severe acute illness with or without fever</li> <li>Moderate or severe acute illness with or without fever</li> </ul>		/ /				
DT, Td	<ul> <li>DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, Tdap</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Age &lt;6 weeks</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> </ul>	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy</li> <li>Moderate or severe acute illness with or without fever</li> <li>Moderate or severe acute illness with or without</li> </ul>						
<ul> <li>DT, Td</li> <li>Hib</li> <li>Hep A</li> </ul>	<ul> <li>DTaP/Tdap only: Encephalopathy (e.g., coma, decreased level of consciousness, prolonged seizures), not attributable to another identifiable cause, within 7 days of administration of previous dose of DTP, DTaP, Tdap</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Age &lt;6 weeks</li> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> </ul>	<ul> <li>History of Arthus-type hypersensitivity reactions after a previous dose of diphtheria-toxoid-containing or tetanus-toxoid-containing vaccine</li> <li>Moderate or severe acute illness with or without fever</li> <li>DTaP/Tdap only: Progressive or unstable neurologic disorder, including infantile spasms, uncontrolled epilepsy, progressive encephalopathy</li> <li>Moderate or severe acute illness with or without fever</li> <li>Moderate or severe acute illness with or without fever</li> </ul>		/ /				

\*<u>https://health.hawaii.gov/docd/files/2019/08/HAR11-157\_EXHIBIT\_B.pdf</u>.

Section 2: For Health Care Provider Use ONLY (MD, DO, ND, APRN-Rx, PA):						
VACCINE	<b>CONTRAINDICATIONS*</b> (Check all that apply to this Patient):	PRECAUTIONS* (Check all that apply to this patient)	FROM:	To:		
□ НРV	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	<ul> <li>Pregnancy</li> <li>Moderate or severe acute illness with or without fever</li> </ul>	/ /	/ /		
□ MMR	<ul> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Pregnancy</li> <li>Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or patients with HIV infection who are severely immunocompromised)</li> <li>Family history of altered immunocompetence</li> </ul>	<ul> <li>Recent (&lt;11 months) receipt of antibody- containing blood product</li> <li>History of thrombocytopenia or thrombocytopenic purpura</li> <li>Need for tuberculin skin testing or interferon- gamma release assay (IGRA) testing</li> <li>Moderate or severe acute illness with or without fever</li> </ul>		/ /		
🗆 мсv	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	Moderate or severe acute illness with or without fever	/ /	/ /		
D PCV	Severe allergic reaction (e.g., anaphylaxis) after a previous dose of PCV13 or any diphtheria-toxoid- containing vaccine or to a component of a vaccine (PCV13 or any diphtheria-toxoid-containing vaccine)	Moderate or severe acute illness with or without fever	/ /	/ /		
□ IPV	Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component	<ul> <li>Pregnancy</li> <li>Moderate or severe acute illness with or without fever</li> </ul>	/ /	/ /		
□ Varicella	<ul> <li>Severe allergic reaction (e.g., anaphylaxis) after a previous dose or to a vaccine component</li> <li>Known severe immunodeficiency (e.g., from hematologic and solid tumors, receipt of chemotherapy, congenital immunodeficiency, long-term immunosuppressive therapy or patients with HIV infection who are severely immunocompromised)</li> <li>Pregnancy</li> <li>Family history of altered immunocompetence</li> </ul>	<ul> <li>Recent (&lt;11 months) receipt of antibody-containing blood product</li> <li>Moderate or severe acute illness with or without fever</li> <li>Receipt of specific antiviral drugs (acyclovir, famciclovir, or valacyclovir) 24 hours before vaccination</li> <li>Use of aspirin or aspirin-containing products</li> </ul>	/ /	/ /		
I certify that in my medical judgement, due to the contraindication(s)/precaution(s) noted above, this student is exempt from the specific vaccine(s) named for						
the period indicated. Health care provider's name/Title (Please Print): License number: Address:						
Health care provider's signature: Date:						

DTaP=Diphtheria, Tetanus, acellular Pertussis, Tdap=Tetanus, diphtheria, acellular pertussis, DT=diphtheria, tetanus, Td=tetanus, diphtheria, Hib=*Haemophilus influenzae* type B, Hep A=hepatitis A, Hep B=hepatitis B, HPV=human papillomavirus, MMR=measles, mumps, rubella, MCV=meningococcal conjugate vaccine, PCV=pneumococcal conjugate vaccine, IPV=inactivated poliovirus vaccine